

Know Your 2017

MALTING BARLEY VARIETIES



Two-Rows

AAC Synergy	(2015)
ABI Voyager	(2014)
AC Metcalfe	(2005)
CDC Copeland	(2007)
CDC Meredith	(2013)
Charles*	(2009)
Conlon	(2000)
Conrad	(2007)
Endeavor*	(2015)
Expedition	(2013)
Harrington	(1989)
Hockett	(2010)
LCS Genie	(2017)
Merit 57	(2010)
Moravian 37	(2010)
Moravian 69	(2010)
ND Genesis	(2016)
Pinnacle	(2011)
Scarlett	(2008)
Wintmalt*	(2013)

*Winter

Six-Rows

Celebration	(2011)
Innovation	(2014)
Lacey	(2000)
Legacy	(2001)
Quest	(2011)
Stellar-ND	(2006)
Thoroughbred*	(2015)
Tradition	(2004)

Variety name & year first recommended


These malting varieties listed in alphabetical order are recommended by AMBA for planting in 2017. When delivered to market in pure carlots of sound, bright, plump, low moisture barley in an acceptable protein range, they may command premium prices over feed barley. Growers are encouraged to contact their local elevator, grain handler or processor to gauge market demand for any variety grown in their region prior to seeding. 2016 crop plantings by variety are included at the end of this publication.

prepared and distributed by

AMERICAN MALTING BARLEY ASSOCIATION, INC.

740 N. Plankinton Avenue, Suite 830, Milwaukee, WI 53203

<http://www.AMBAinc.org>



	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>
Six-Rows	(% of Six-Row Malting Varieties)			
CELEBRATION	4.4%	2.8%	0.1%	0.1%
INNOVATION	1.6%	2.6%	2.2%	4.1%
LACEY	33.5%	18.3%	24.6%	27.7%
LEGACY	1.1%	1.3%	1.0%	1.2%
QUEST	0.5%	3.0%	2.1%	*
STELLAR-ND	0.1%	*	*	*
TRADITION	58.2%	71.8%	69.8%	67.0%

	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>
Two-Rows	(% of Two-Row Malting Varieties)			
AAC SYNERGY	*	*	*	0.7%
ABI VOYAGER	2.3%	5.0%	10.7%	15.5%
AC METCALFE	26.0%	23.3%	22.7%	16.3%
BC 100	*	*	*	0.3%
CDC COPELAND	5.4%	6.9%	6.3%	9.3%
CDC MEREDITH	0.9%	2.5%	1.7%	0.4%
CHARLES	0.8%	0.9%	0.5%	0.9%
CONRAD	18.8%	18.0%	10.2%	9.6%
EXPEDITION	0.3%	0.3%	0.5%	0.8%
FULL PINT	*	*	0.1%	0.1%
HARRINGTON	1.8%	2.1%	1.4%	0.9%
HOCKETT	6.7%	6.0%	15.2%	10.3%
MERIT	0.5%	0.4%	*	*
MERIT 57	9.3%	7.9%	6.5%	6.1%
MORAVIAN 37	*	*	*	*
MORAVIAN 69	14.9%	16.9%	11.7%	14.8%
MORAVIAN 115	8.4%	4.8%	7.2%	8.1%
ND GENESIS	*	*	0.3%	1.6%
PINNACLE	3.4%	3.8%	4.3%	1.3%
SCARLETT	0.3%	0.4%	0.4%	0.3%
WINTMALT	*	0.2%	0.2%	0.4%

* Less than 0.1%

Source: AMBA Industry Barley Variety Surveys (CO, ID, MN, MT, ND, WA, WY)

VARIETY IDENTIFICATION PROCEDURE


VARIETY IDENTIFICATION IS NOT DIFFICULT

Surveys show that in any one community there are usually not more than five or six varieties. With careful study of typical, pure kernel samples one can learn to recognize local varieties with a high degree of accuracy.

GUIDES IN IDENTIFYING KERNELS OF BARLEY VARIETIES


1. Study type samples - look at the sample as a whole, not at only a few individual kernels. Line up a row of 10-20 kernels pointing the base end toward the light source.
2. Use a magnifier with a good strong light - north daylight is best.
3. Know which varieties are being grown in your area - keep an eye open for any new introductions. Study varietal kernel characteristics for each new crop.

Do not be afraid to look and look, and look again.



TO IDENTIFY A SAMPLE

- A. Check general appearance of the sample. A pure sample will usually appear uniform. A sample containing a mixture of varieties generally has a non-uniform appearance.
- B. Determine the major kernel characters as follows:



1. Six-Row or Two-Row - The lateral or side kernels in six-row varieties are twisted. Therefore, in six-row varieties, two-thirds of the kernels are twisted. In two-row varieties, which have no lateral kernels, all kernels are straight and symmetrical and none are twisted.
2. White or Blue pearl (aleurone) – This is an easy character to determine with the use of a barley pearling machine. Sometimes blue varieties may have very weakly developed blue color and can be confused with white pearls.
3. Rough or Smooth Beards (awns) – There are usually enough broken pieces of beard or short pieces of beard attached to the kernel so that this character can easily be determined. Most smooth or semi-smooth bearded varieties have a few barbs at the tip but have no barbs on the rest of the beard.

4. Long or Short Hairs on Rachilla - This is an excellent character to use but a hand magnifying lens is needed to see the rachilla hairs. The difference between long and short hairs is very pronounced and does not vary from year to year or in different areas. Some varieties have aborted rachillas, these can be seen readily with the naked eye.

C. If the above characters are not sufficient to identify a variety, other characters will have to be used. The most useful are described below:

1. The basal mark can vary from a depression to a transverse crease. A depression is a smooth uniform hollow or “dimple”. A transverse crease is a sharp fold or line.
2. Crease shape may vary from wide, open and V-shaped from base of kernel to narrow and almost closed. In some varieties the crease is closed in the bottom half of the kernel and flaring at the tip end. This character will vary with growing conditions and plumpness of kernel.

3. Barbs on lateral veins can be seen with a magnifying glass or felt with the finger. Their number varies from none to many in different varieties. These barbs are not related to the barbs on the beards of rough bearded varieties.

4. Hairs on glumes require careful observation with a hand magnifying lens to determine the length of hairs and their location on the glume. This can be a very useful character in varieties such as Foster where the glumes are covered with long hairs.

5. Kernel shape varies with growing conditions but can be of value in a local area. The relationship between length and width and fullness in different parts of the kernel are useful characters. Kernel shape may be smooth and tapered or there may be prominent bulges or other distinctive features.

6. Wrinkling of the hull varies from very fine to coarse. Many two-row varieties have numerous, very fine wrinkles. Some varieties, such as Steptoe, have very little wrinkling and may have a smooth hull.

C. In any area one may find additional characters which will be of use in identifying local varieties. It is important to study locally grown samples each year as some kernel characters may vary depending on the season and location at which a variety is grown.

RACHILLA HAIRS



Rachilla hairs: left – long, center – short, right - aborted

KERNEL CHARACTERS



Hairiness of glumes: left to right – covered, in band, on midline, without hairs or smooth.

GLUME HAIRS



Variations in

width.

crease shape and

CREASE SHAPE

KERNEL CHARACTERS



Basal mark: left – depression, center – depression tending to crease, right – transverse crease.

BASAL MARK




Wrinkling of hulls: left – slightly wrinkled, center – semi-wrinkled, right – wrinkled.

HULL WRINKLING

Barley Variety Survey - 2016

PERCENT OF ACREAGE BY STATE




Statistics obtained from companies contracting malting barley in the indicated states. While the majority of the contracted acreage is included, there may be some acres not covered by this survey. Open market malting barley, food and feed varieties are included in "unknown". Varieties with acreage equal to or less than 0.10% of a state's acreage are reported as "other malting".



Barley Variety Survey - 2016

PERCENT OF ACREAGE BY CROP REPORTING DISTRICTS



REPORT BASED UPON INDUSTRY SURVEY OF ACRES CONTRACTED BY VARIETY. PERCENTAGES BASED ON TOTAL DISTRICT ACRES REPORTED BY THE USDA/NASS. DISTRICT (OR COMBINED DISTRICTS) ACREAGE IN 2016 IS INDICATED BENEATH THE DISTRICT LIST OF VARIETIES.

¹Unknown represents acreage of malting, feed, and food barley varieties not reported to AMBA in the survey.
²Oth & Unk stands for Other and Unknown where Other represents contracted area of minor or low acreage varieties in a region. Unknown is explained above.

